

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A61K 38/16		A1	(11) International Publication Number: WO 96/32129 (43) International Publication Date: 17 October 1996 (17.10.96)
(21) International Application Number: PCT/US96/04674 (22) International Filing Date: 10 April 1996 (10.04.96) (30) Priority Data: 08/419,066 10 April 1995 (10.04.95) US (60) Parent Application or Grant (63) Related by Continuation US 08/419,066 (CIP) Filed on 10 April 1995 (10.04.95) (71) Applicant (for all designated States except US): KANSAS STATE UNIVERSITY RESEARCH FOUNDATION [US/US]; 146 Durland Hall, Manhattan, KS 66506 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): BLECHA, Frank [US/US]; 2161 Deep Creek Road, Manhattan, KS 66502 (US). SHI, Jishu [US/US]; 400 Jardine Terrace, S-23, Manhattan, KS 66502 (US). (74) Agent: COLLINS, John, M.; Hovey, Williams, Timmons & Collins, Suite 400, 2405 Grand Boulevard, Kansas City, MO 64108 (US).			(81) Designated States: AL, AM, AT, AU, AZ, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IS, JP, KE, KG, KP, KR, KZ, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, US, UZ, VN, ARIPO patent (KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
(54) Title: SYNTHETIC PEPTIDES THAT INHIBIT LEUKOCYTE SUPEROXIDE ANION PRODUCTION AND/OR ATTRACT LEUKOCYTES (57) Abstract Methods of inhibiting leukocyte O ₂ ⁻ production and attracting leukocytes using specific peptides are disclosed. These peptides include the proline-arginine (PR)-rich antimicrobial peptide known as PR-39 and truncated analogs thereof. These peptides can be used as medicaments that fight infection by attracting leukocytes to a wound site, yet restrict tissue damage at the wound site caused by excessive oxygen radicals produced by these leukocytes.			

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

-1-

SYNTHETIC PEPTIDES THAT INHIBIT LEUKOCYTE SUPEROXIDE
ANION PRODUCTION AND/OR ATTRACT LEUKOCYTES

5 Background of the Invention

1. Field of the Invention

The present invention is broadly concerned with methods of inhibiting leukocyte superoxide anion (O_2^-) production and methods of attracting a leukocyte to a location using peptides. More particularly, the invention relates to methods employing a naturally occurring proline-arginine (PR)-rich antimicrobial peptide known as PR-39 and analogs thereof; these peptides can be used as medicaments that fight infection by attracting leukocytes to a wound site, yet restrict tissue damage at the wound site caused by excessive oxygen radicals produced by these leukocytes.

15 2. Description of the Prior Art

Infectious diseases are a primary cause of morbidity and mortality in humans and animals. These maladies range from the troublesome, e.g., *Escherichia coli* diarrhea which is caused by the consumption of contaminated food and drinks, to the deadly, e.g., AIDS.

20 A variety of antimicrobial agents have been developed to combat infectious diseases. Recently, several types of antimicrobial peptides have been discovered. Such peptide antimicrobials are produced by many biological organisms and are important components of host defense mechanisms. (Boman, 1991; Zasloff, 1992; Gabay, 1994; Boman, 1995, Martin et al., 1995)
25 For example, defensins are expressed in several mammalian species (Lehrer et al., 1993), magainins have been identified in the skin and intestine of frogs (Zasloff, 1987; Moore et al., 1992), and cecropins have been isolated from insects and pigs (Steiner et al., 1981; Lee et al., 1989). These natural antimicrobials are lytic peptides that kill microorganisms by pore-forming,
30 membrane-damaging mechanisms (Boman et al., 1993; Maloy et al., 1995).

The patent art also discloses antimicrobial peptides. U.S. Patent No. 5,234,716 describes broad spectrum tryptophan antimicrobial peptides, and U.S. Patent No. 5,202,420 discloses tracheal antimicrobial peptides.

35 Recently, a group of PR-rich antibacterial peptides have been identified. Bactenecins 5 and 7 have been isolated from bovine neutrophils

-2-

(Gennaro et al., 1989; Litteri et al., 1993), and PR-39 was first isolated from the porcine small intestine (Agerberth et al., 1991) and identified recently in porcine and human neutrophils (Shi et al., 1994b; Shi et al., 1995). Although these PR-rich antibacterial peptides share a similar high content of proline (47, 47, and 49%, respectively) and arginine (21, 29, and 26%, respectively), they possess different killing mechanisms.

Similar to other lytic peptides, bactenecins kill bacteria by a membrane-permeability-associated mechanism (Maloy et al., 1995); however, PR-39 was found to kill bacteria by interfering with DNA and/or protein synthesis (Boman et al., 1993). Furthermore, PR-39 has been isolated from wound fluid and was shown to induce syndecan expression on mesenchymal cells (Gallo, 1994). Because syndecans are important in wound repair, this finding suggests that PR-39, in addition to its antibacterial properties, may have a larger role in inflammatory processes and tissue repair.

Neutrophils represent a first line of defense against infections; they are the first white blood cells to arrive at sites of infection and are well-equipped to sequester and eliminate pathogens. These cells possess multiple antimicrobial defense mechanisms, including both oxidative and nonoxidative microbial killing processes (Klebanoff, 1992; Selsted et al., 1995). Nonoxidative neutrophil defense mechanisms include several antibacterial peptides including PR-39. Phagocyte oxidative defense mechanisms are initiated by a plasma membrane-bound enzyme complex called reduced nicotinamide dinucleotide phosphate (NADPH) oxidase (Rotrosen, 1992). This multicomponent enzyme catalyzes the reduction of molecular oxygen to O_2^- using NADPH as an electron donor. Although O_2^- and other reactive oxygen intermediates are important components of host defense, these highly toxic oxidants also cause significant tissue injury in inflammatory diseases and ischemia-reperfusion injury (Shasby et al., 1982; Malech, 1987; Demling, 1990; Martinez-Cayuela, 1995; Granger et al., 1995). Thus, their generation and inactivation must be tightly regulated.

At least five proteins comprise the NADPH oxidase complex; a membrane flavocytochrome b_{558} , which is composed of two subunits (gp91^{phox} and p22^{phox}); and three cytosolic components (p47^{phox}, p67^{phox}, and a GTP-binding protein named p21^{rac}) (Rotrosen et al., 1992; Abo et al., 1992). Although mechanisms for activation and assembly of NADPH oxidase have not

-3-

been elucidated fully, it is clear that multiple protein-protein interactions among its components are regulated by a number of signaling intermediates (McPhail et al., 1993). The assembly of phagocyte NADPH oxidase requires protein-protein interactions between Src homology 3 (SH3) domains in cytosolic components and proline-rich regions in other components (Leto et al., 1994; Sumimoto et al., 1994; Finan et al., McPhail, 1994; de Mendez, 1996).

Summary of the Invention

The present invention is predicated upon the discovery that specific peptides (e.g., PR-39) are capable of 1) inhibiting O_2^- synthesis by leukocyte enzymes (e.g., NADPH oxidase), and 2) attracting leukocytes (e.g., neutrophils). These peptides can be used as novel medicaments that fight infection by attracting leukocytes to a wound site, yet restrict tissue damage at the wound site caused by excessive oxygen radicals produced by these leukocytes. Preferably, these peptides have a sequence included in PR-39 (e.g., Sequence ID Nos. 1 and 2 for peptides capable of inhibiting O_2^- production, and Sequences ID Nos. 1, 2, 5, 6, and 7 for peptides capable of attracting leukocytes). Advantageously, these peptides are synthesized and have lengths of less than 60 amino acid residues, and the leukocytes upon which the peptides act are mammalian (e.g., porcine) leukocytes. This invention was made with government support under Grant 93-37206-9351 awarded by the United States Department of Agriculture. The government has certain rights in the invention.

Brief Description of the Drawings

Figure 1 is the amino acid sequence of PR-39 and various truncated analogs thereof, wherein the single letter amino acid code is used and the left side of the sequence is the NH_2 -terminal end;

Fig. 2 is a protein hydrophilicity (solid line) and antigenicity (dashed line) plot for PR-39 wherein the hydrophilic and antigenic domains are above the horizontal line and the hydrophobic domains are below the horizontal line;

Fig. 3 is a photograph of an AU-PAGE analysis of synthetic peptides wherein each lane contains ten micrograms of peptide; lane 1 = PR-14; lane 2 = PR-15; lane 3 = PR-16; lane 4 = PR-26; lane 5 = PR-39; and lane

-4-

6 = neutrophil granule extract, with calculated molecular weights (MW_{cal}) over the experimental mass spectrometry data (MW_{exp}) listed above each lane for the respective synthetic peptides;

5 Fig. 4 is a photograph of the gel shown in Fig. 3 after being subjected to a gel-overlay assay, wherein the clear zones indicate antibacterial activity;

10 Fig. 5 is a photograph of an agar plate showing the results of an antibacterial "lawn-spotting" assay using *Salmonella typhimurium* wherein the spot positions are numbered on the left of the photograph; position 1 = medium control (0.01% acetic acid); positions 2, 3, and 4 = PR-14, PR-15, and PR-16, respectively, at 1 mg/ml in 0.01% acetic acid; positions 5 and 6 = 100 μ moles and 50 μ moles, respectively, of PR-26 in phosphate buffer saline, pH 7.4; positions 7, 8, 9, and 10 = 100, 50, 25, and 10 μ moles, respectively, of PR-26 in 0.01% acetic acid; and positions 11, 12, and 13 = 100 μ moles, 50 μ moles, and 25 μ moles, respectively, of PR-39 in 0.01% acetic acid;

15 Fig. 6 is a photograph of a 96-well microtiter plate used to determine the minimal inhibitory concentrations of PR-26 and PR-39 against enteric bacteria;

20 Fig. 7 is a graph illustrating the postantibiotic effect of PR-26 and PR-39 against *S. typhimurium*;

Fig. 8 is a graph illustrating the phagocytic susceptibility of *S. choleraesuis* after exposure to PR-26 and PR-39;

Fig. 9 is a graph illustrating the chemotaxis of porcine neutrophils by PR-39 and analogs thereof;

25 Fig. 10 is a graph illustrating a dose response of PR-39 for neutrophil chemotaxis;

Fig. 11 is a graph illustrating the cytotoxicity of PR-26 and PR-39 in intestinal epithelial cells;

30 Fig. 12 is a graph illustrating percent survival of A/J mice 15 days after an oral challenge with *S. typhimurium* followed with 0, 100, or 250 μ g PR-26;

Fig. 13 is a graph illustrating survival of mice infected with *S. typhimurium* intraperitoneally (i.p.) followed with 0, 50, or 100 μ g PR-26;

-5-

Fig. 14 is a graph illustrating survival of mice infected with *S. typhimurium* i.p. followed with 50 µg PR-26 at time 0, 24, 48, or 72 hours post-infection;

Fig. 15 is a graph illustrating that inhibition of whole-cell NADPH oxidase activity requires preincubation with PR-39;

5 Fig. 16 is a graph illustrating whole-cell NADPH oxidase inhibition by PR-39 and analogs thereof;

Fig. 17 is a graph illustrating cell-free NADPH oxidase inhibition by PR-39 and analogs thereof;

10 Fig. 18 is a photograph of blots illustrating the binding of biotinylated PR-39 to a 47 kDa neutrophil cytosol protein;

Fig. 19 is a photograph of a blot and a corresponding graph illustrating the binding of PR-39 to recombinant p47^{phox};

15 Fig. 20 is a photograph of blots illustrating the degrees to which PR-26 and PR-39 block the interaction between GST-p47^{phox} and recombinant p22^{phox};

Fig. 21 is a photograph of a blot illustrating the degrees to which PR-19 and PR-23 block the interaction between GST-p47^{phox} and recombinant p22^{phox}; and

20 Fig. 22 is a photograph of a blot illustrating the degrees to which PR-14, PR-15, PR-16, and PR-26 block the interaction between GST-p47^{phox} and recombinant p22^{phox}.

Detailed Description of the Preferred Embodiment

25 The following example illustrates the preferred practice of the invention. It is to be understood, however, that this example is provided by way of illustration only and nothing therein should be taken as a limitation upon the overall scope of the invention.

Example

30 MATERIALS AND METHODS

Peptide design and synthesis. Theoretical predictions of peptide characteristics, relative to hydrophilicity, hydrophobicity, and antigenicity, were accomplished using a computer software program (Peptide Companion, Peptide International, Louisville, KY). The following peptides were synthesized:

35 PR-39, PR-26, PR-23, PR-19, PR-16, PR-15, and PR-14; the sequences of

these peptides appear in the Sequence Listing as Sequence ID Nos. 1 through 7, respectively. Peptides were synthesized by the solid-phase method using t-Boc chemistry with an Applied Biosystems Model 431 Peptide Synthesizer (ABI, Foster City, CA). Amino acid derivatives having the L-configuration were used. Peptide purification and characterization were conducted as described previously (Shi et al., 1994b; Shi et al., 1996). Briefly, the peptides were purified on a reversed-phase high-performance liquid chromatography (RP-HPLC) system (Beckman Instruments, Fullerton, CA) with a Vydac 218 TP C₁₈ column (0.46 x 25 cm), analyzed by fast-atom mass spectrometry (AUTOSPEC-Q; VG Analytical Ltd., Manchester, United Kingdom), and visualized by acid-urea polyacrylamide gel electrophoresis (AU-PAGE). In the AU-PAGE analysis, peptides (10 µg) were dissolved in 20 µl of sample buffer (3M urea with 5% acetic acid) and the gels were run at 150V for approximately 15 min. or until the dye front (methyl green) had migrated to the end of the gel. The gel was stained with 0.3% amido black.

Antibacterial activity assays. Synthetic peptides were evaluated for antibacterial activity by previously described gel-overlay and "lawn-spotting" assays (Shi et al., 1994b), by determination of the minimal inhibitory concentrations (MICs) and the minimal bactericidal concentrations (MBCs) of PR-26 and PR-39, and by determination of the postantibiotic effect (PAE) of PR-26 and PR-39 and the bacterial susceptibility to neutrophil phagocytosis by PR-26 and PR-39.

(i) Gel-overlay assay. Peptides were subjected to AU-PAGE as described previously (Shi et al., 1994b). The acid-urea gels were overlaid with 10⁶ bacteria (*E. coli*, ATCC 25922) in 3% trypticase soy broth (TSB) and 1% agarose medium, and the overlaid gels then were incubated at 37°C for 18 hr. Bactericidal activity was indicated by clear zones on the agarose gels.

(ii) Lawn-spotting assay. Lawns of bacteria (*E. coli*, ATCC 25922, or *S. typhimurium*, KSU isolate 7) were made on sheep-blood or brain-heart-infusion agar plates. After drying at room temperature for 10 min, 5 µl of the various peptides (dissolved in 0.01% acetic acid or phosphate buffered saline (PBS), pH 7.4) and a medium control were spotted on the surface of the bacterial lawn. Plates were incubated at 37°C for 18 hr. Bactericidal activity was indicated by a clear zone on the bacterial lawn (Shi et al., 1994b).

-7-

(iii) MICs and MBCs. For peptides that showed antibacterial activity using the gel-overlay and lawn spotting assays, MICs and MBCs were determined by the microdilution broth method (NCCLS, 1990). Briefly, 50 µl of twofold serial dilutions (128 to 0.25 µM) of synthetic peptides were dispensed into wells of 96-well tissue culture plates. Bacteria (*E. coli*, ATCC 25922; *E. coli*, K88; *S. typhimurium*, fresh isolate from a dog; *Salmonella choleraesuis* ATCC 6962; *Streptococcus suis*, fresh isolate from porcine spleen; and *Staphylococcus aureus*) from sheep-blood or brain-heart-infusion agar plates were standardized to 0.5 McFarland in demineralized water using a radiometer/sensititre (Chelsea Instrument Ltd., England). The water/bacteria suspension (100 µl) was immediately transferred to 10 ml of cation-adjusted Muller-Hinton broth and 50 µl of the bacterial suspension then were added to each well of the microtiter plate. Plates were incubated for 20 hr. at 37°C and MICs were determined. Ten microliters of bacterial suspensions in Muller-Hinton broth also were diluted to determine the actual bacterial concentration using standard colony forming unit (CFU) counting. After determination of the MICs, 10 µl of each bacteria-peptide suspension were plated on sheep-blood or brain-heart-infusion agar plates and incubated for 24 hr. at 37°C to determine the MBCs. MBC was considered that peptide concentration that inhibited 99.9% of the original CFUs (NCCLS, 1990).

(iv) PAE. *S. typhimurium* was used to evaluate the PAE of PR-26 and PR-39. Stationary phase bacteria were adjusted to 5×10^7 bacteria/ml in brain-heart-infusion agar and incubated with different concentrations of PR-26 or PR-39 at 37°C for 2 hr. Control tubes without PR-peptides were treated in an identical manner to the experimental tubes. PR-26 and PR-39 were removed by centrifugation (13,600 x g for 1 min.) and 100 µl of the bacteria were resuspended in 0.9 ml of PR-peptide-free brain-heart-infusion agar and incubated at 37°C. Bacteria (20 µl) were diluted in sterile saline immediately after removal of the PR-peptides and then at hourly intervals, and 20 µl aliquots were spread on nutrient agar plates. Viable bacteria were counted. Tests were repeated on three different days. PAE was determined by calculating the difference in time required for the number of test and control bacteria to increase 1 log₁₀ above the number present immediately after removal of PR-peptides from the test cultures. The results were expressed as the mean ±

-8-

standard deviation. A PAE greater than 30 min. was considered significant (MacKenzie and Gould, 1993).

(v) Susceptibility to neutrophil phagocytosis. Porcine neutrophils were isolated from 6 to 8 week-old-crossbred pigs by density-gradient centrifugation and hypotonic lysis as previously described (Shi et al., 1994a). *S. choleraesuis*, ATCC 6962 was used in this experiment. Bacteria were incubated with PR-26 or PR-39 for 10 min. at 37°C. Peptides were removed from the bacterial cultures by centrifugation at 13,600 x g for 1 min. Bacteria were resuspended in 1 ml of PBS and 0.1 ml of bacteria was mixed with 2 x 10⁶ porcine neutrophils. The final volume was adjusted to 0.5 ml, 15% porcine serum. Bacteria without neutrophils and neutrophils without bacteria were used as controls. Tubes were incubated at 37°C in a reciprocating water bath at 110 oscillations/min. Aliquots of 50 µl from the experimental tubes were used to prepare slides using a Cytospin 2 centrifuge (Shandon Products Ltd., Pittsburgh, PA). Slides were stained with LeukoStat solutions (Fisher, Pittsburgh, PA). Phagocytosis was determined by light microscopy at a magnification of 1,000. At least 200 neutrophils were examined. The degree of phagocytosis was calculated according to the following formula: phagocytic index = (percentage of neutrophils containing at least one bacteria) x (mean number of bacteria per positive cell). Tests were repeated on three different days. Results were expressed as the mean ± standard deviation.

Influence of PR-39 on neutrophil chemotaxis. Chemotaxis of porcine neutrophils was measured by the procedure of Salak et al. (1993). Briefly, PR-14, PR-15, PR-16, PR-26, or PR-39 (30 µl in Dulbecco's Modified Eagle's medium) were placed in the bottom chamber of a modified Boyden chamber (Neuro Probe, Cabin John, MD) and porcine neutrophils (50 µl at 5 x 10⁶ cells/ml) were placed in the top chamber. The chambers were incubated at 37°C for 30 min. Cells that migrated through the porous membrane (pore size 5 µm) were stained using LeukoStat solution and enumerated. Five microscope fields were counted and the cells that migrated through the membrane were standardized to the medium control and referred to as the migration index.

Influence of PR-26 and PR-39 on intestinal epithelial cells. A nonradioactive assay based on the cellular conversion by viable cells of a tetrazolium salt into a blue formazan product was used to determine if PR-26

-9-

or PR-39 were toxic to intestinal epithelial cells. Ninety-six-well microtiter plates were seeded with the rat intestinal epithelial cell line, IEC-6, (5×10^4 cells/well) in DMEM containing 10% fetal bovine serum, 1% antibiotic/antimycotic, and 0.1 bovine insulin, and incubated at 37°C for three days to achieve confluency. Cells then were incubated with different concentrations of PR-26 or PR-39 in medium without antibiotic/antimycotic for three days. Well contents then were aspirated and monolayers were washed with medium. Medium (100 µl) and 15 µl of dye solution (Promega, Madison, WI) were added to each well and plates were incubated for 4 hr. at 37°C. Solubilization buffer (100 µl, Promega, Madison, WI) was added to each well and plates were incubated overnight to allow solubilization of the formazan crystals. Absorbance then was read at 570 nm using a microplate reader.

Effect of PR-26 on survival of mice challenged with *S. typhimurium*.

Three separate *S. typhimurium* challenge studies were conducted with mice. In the first experiment, 45 A/J mice (Jackson Labs, Bar Harbor, ME) were orally challenged with *S. typhimurium*. Feed was withdrawn 4 hrs. prior to an oral inoculation. Twenty min. prior to the inoculation with *S. typhimurium* mice were given a 30 µl dose of a 10% sodium bicarbonate solution in PBS. *S. typhimurium*, strain KSU007, was started in brain heart infusion broth overnight and then grown for 6 hrs. in Luria Broth Base (GIBCO, Gaithersburg, MD). After centrifugation, bacteria were suspended in a 1% gelatin solution (final concentration 1.24×10^{10} *S. typhimurium*/ml), and then 25 µl were given orally. Immediately following oral inoculation with *S. typhimurium*, mice were given 0, 100, or 250 µg of PR-26 in PBS. A second dose of PR-26 or PBS was given 4 hrs. later.

In the second experiment, 45 Balb/C (*Salmonella* susceptible) mice were given i.p. injections of *S. typhimurium* (2×10^4 /ml) that had been grown as previously described, but resuspended in PBS. Mice then received either 0, 50, or 100 µg of PR-26 in 100 µl PBS i.p. immediately after the *S. typhimurium*.

In the third experiment, the efficacy of timing of delivery of PR-26 was evaluated. The protocol was the same as in the second experiment, except that only the 50 µg/mouse dose of PR-26 was used and the PR-26 was delivered immediately (time 0), or 24, 48, or 72 hrs. after the *S. typhimurium*

-10-

injection. A control group received PBS at time 0. Fifteen mice were used in all treatments except the 72-hour treatment, which had 16.

Regulation of neutrophil O_2^- production by PR-26 and PR-39.

5 (i) O_2^- production assays. Whole-cell O_2^- production by porcine peripheral blood neutrophils was determined by the superoxide dismutase-inhibitable reduction of ferricytochrome c as previously described (Shi et al., 1994a). Cell-free O_2^- production was measured as previously described (Leto et al., 1991) using 96-well plates and a Molecular Devices Thermomax microplate reader (Menlo Park, CA). Reactions (100 μ l) contained 10^6 cell
10 equivalents of human neutrophil cytosol and 5×10^5 cell equivalents of deoxycholate-solubilized membranes prepared from human peripheral blood neutrophils. Reaction mixtures contained 50 mM potassium phosphate (pH 7), 0.2 mM acetylated cytochrome c, 4 mM $MgCl_2$, 1 mM EGTA, 10 μ M FAD, 1 μ M GTP- γ S, and 200 μ M NADPH. The reactions were initiated by addition of 40
15 μ M arachidonic acid. Control reactions contained 5 μ g of superoxide dismutase. Superoxide anion generation was calculated based on superoxide dismutase-inhibitable changes in cytochrome c absorbance observed at 551 nm. The reactions were followed for 20 min. after addition of arachidonic acid, with absorbance readings taken at 1-min. intervals. Maximum rates of
20 superoxide generation were calculated from a linear least squares fit of 10 consecutive 1-min. data points. Determinations were based on reactions performed in duplicate.

(ii) Binding assays. Neutrophils were disrupted by sonication, and unbroken cells, debris, and granules were removed by centrifugation
25 (13,000 x g for 60 min.). Cytosol fractions were prepared by centrifugation (100,000 g for 45 min.) of the supernatants to remove residual membranes. Cytosolic proteins (15 μ g in reducing buffer) were separated on a 7.5% SDS PAGE gel, transferred to PVDF membranes, and probed with biotinylated PR-39 (0.3 μ M) for 2 hr. Binding between PR-39 and cytosolic proteins were
30 visualized by a chemiluminescent streptavidin alkaline phosphatase system using an Epson scanner (Torrance, CA) and the NIH *Image* program. PR-39 was labeled with biotin hydrazide at the carboxy-terminus (Pierce Chemical Co., Rockford, IL) and purified by gel filtration chromatography.

Competitive ligand-blot assays were performed by incubating
35 recombinant p47^{phox} (0.02 μ M) with or without various concentrations (0.006,

0.06, and 0.6 μ M) of PR-39 for 1 hr. at 37°C and then with biotinylated PR-39 (0.06 μ M) for an additional 1 hr. at 37°C. The solutions were dialyzed against ddH₂O for 1 hr. at 4°C before blotting to nitrocellulose (5 μ l) and detection with a streptavidin alkaline-phosphatase detection system as described above.

5 Methods for production of the glutathione S-transferase (GST)-p47^{phox}-SH3 (residues 151-284) fusion protein have been previously described (Leto et al., 1994). For solution binding assays, lysates of recombinant p22^{phox} protein (residues 127-195) from baculovirus-infected cells (2.5 x 10⁴ cell equivalents per assay) were mixed with peptides and incubated for 1 hr. at
10 20°C with 100 μ l of GST-p47^{phox} bound to glutathione-Sepharose beads (15% suspension). Bound proteins were washed three times in 15 volumes of ice-cold 100 mM KCl/3 mM NaCl/3.5 mM MgCl₂/0.15 mM phenylmethanesulfonyl fluoride/10 mM Pipes (pH 7.5), eluted with 1% SDS, and analyzed by SDS-PAGE followed by immunoblotting with mouse anti-p22^{phox} antibody (mAb 449;
15 A. Verhoeven, Central Laboratories of the Netherlands Red Cross). Controls included blotting of recombinant p22^{phox} lysate (5 x 10³ cell equivalents) alone (p22) and bound complexes detected without competing PR peptides (-).

RESULTS AND DISCUSSION

20 Peptide design and synthesis. PR-39 and six analogs were synthesized and named PR-39 (whole molecule), PR-26 (NH₂-terminal segment 1 to 26), PR-23 (control segment 4 to 26), PR-19 (NH₂-terminal segment 1 to 19), PR-16 (central segment 11 to 26), PR-15 (COOH-terminal segment 25 to 39), and PR-14 (NH₂-terminal segment 1 to 14). Peptide sequences are illustrated in Fig.
25 1. Rationale for design of the peptides was based on the graphic protein hydrophilicity scale (Fig. 2), since the hydrophilicity profile indicates locations of important interaction sites such as antibody and receptor binding sites (Hopp, 1985). PR-26 was designed to mimic the hydrophilicity profile of PR-39 which has a hydrophilic NH₂-terminus and a hydrophobic COOH-terminus. As
30 most endogenous antibacterial peptides are cationic molecules, PR-14, the highest positively charged segment (43% vs. 25% of whole molecule), was designed to test if cationicity itself was enough for antibacterial activity. The central domain, PR-16, having the average positive charge intensity (25%) was synthesized as the control for PR-14. PR-15, the COOH-terminus of PR-39
35 was designed to determine if it was one of the functional domains of PR-39.

-12-

PR-23 was designed to evaluate the importance of the first three arginine residues of PR-26. PR-19 was designed to assess the contribution of the COOH-terminus to the activity of PR-26.

5 Calculated molecular weights and experimental determinations of the mass of synthetic peptides suggested that these peptides possessed the designed sequences (1813, 1777, 1939, 3230, and 4720 calculated molecular weight vs 1812, 1777, 1941, 3230, and 4720 experimental determination of mass for PR-14, PR-15, PR-16, PR-26, and PR-39, respectively). Synthetic peptides (>95% purity in RP-HPLC) were subjected to AU-PAGE to further
10 determine the purity and charge intensity. Fig. 3 shows the results of this analysis for PR-14, PR-15, PR-16, PR-26, and PR-39 (data not shown for PR-19 and PR-23). As expected, PR-14 migrated in the front since it has the highest positive charge intensity; PR-15, having the lowest positive charge intensity, migrated far behind PR-14 even though its mass is slightly less than
15 PR-14. Because PR-16 and PR-26 are smaller molecules than PR-39, they migrated faster than the parent molecule. Native PR-39 and synthetic PR-39 behaved identically in AU-PAGE and RP-HPLC analyses (data not shown).

Antibacterial activity assays.

(i) Gel-overlay assay. In the gel-overlay bactericidal assay,
20 only PR-26 and PR-39 were found to have antibacterial activity against *E. coli*. Fig. 4 shows the results of this assay for PR-14, PR-15, PR-16, PR-26, and PR-39 (data not shown for PR-19 and PR-23).

(ii) Lawn-spotting assay. In the "lawn-spotting" antibacterial assay, PR-14, PR-15, PR-16, PR-19, PR-23, PR-26, and PR-39, and the
25 combination of PR-14, PR-15, and PR-16, were tested. Fig. 5 shows the results of this assay for PR-14, PR-15, PR-16, PR-26, and PR-39 (data not shown for PR-19, PR-23, and the combination of PR-14, PR-15, and PR-16). Only PR-26 and PR-39 had antibacterial activity. All of the other segments and their mixtures showed no antibacterial activity even at 1 mg/ml.

30 These results suggest that: 1) the very cationic NH₂-terminus of PR-39 is not sufficient for antibacterial activity; 2) the COOH-terminal segment 27 to 39 does not contribute to the antibacterial activity of PR-39; 3) PR-26, the NH₂-terminal segment 1 to 26, contains the antibacterial domain of PR-39; 4) the NH₂-terminal segment 1 to 3 is essential for antibacterial activity; (5) the
35 COOH-terminal segment 20 to 26 is essential for antibacterial activity; and 6)

-13-

certain secondary structure conformation is required for the antibacterial activity of PR-26 and PR-39 since segment mixtures did not have any antibacterial activity. The "lawn-spotting" assay also showed that PR-26 had greater antibacterial activity against *E. coli* and *S. typhimurium* than PR-39 (Fig. 5).

(iii) MICs and MBCs. The MICs of PR-26 and PR-39 for *E. coli*, 25922; *E. coli*, K88; *S. typhimurium*; *S. choleraesuis*; *S. suis*; and *S. aureus* are shown in Fig. 6 and Table 1. For the enteric, Gram-negative bacteria, the MICs of PR-26 and PR-39 ranged from 1 to 4 μ M; PR-26 had a lower MIC than PR-39. Similarly, the MBCs of PR-26 were the same or lower than the MBCs of PR-39 and ranged from 2 to 8 μ M for the Gram-negative bacteria (Table 1). These findings suggest that PR-26 may be an effective antibiotic against enteric, Gram-negative bacteria such as *E. coli* or *Salmonella*.

Table 1. MICs (μ M) and MBCs (μ M) of PR-39 and PR-26 against six strains of bacteria.

Bacteria	MIC		MBC	
	PR-39	PR-26	PR-39	PR-26
<i>Escherichia coli</i> , 25922	4	2	8	8
<i>Escherichia coli</i> , K88	2	1	4	4
<i>Salmonella typhimurium</i>	4	2	4	2
<i>Salmonella choleraesuis</i>	2	1	4	2
<i>Streptococcus suis</i>	>64	16	>64	>64
<i>Staphylococcus aureus</i>	>250	>250	ND*	ND*

*Not determined

(iv) PAE. The relationship between peptide concentration and the duration of PAE of PR-26 and PR-39 is shown in Fig. 7. Suboptimal peptide concentrations (0.1 MIC) only caused a slight growth delay in both PR-26- and PR-39-treated bacteria. However, at 1 and 8 MICs, PAEs against *S. typhimurium* were significantly increased. These findings agree with other

-14-

antibacterial data and show clearly that PR-26 limits the growth of enteric bacteria.

5 (v) Susceptibility to neutrophil phagocytosis. Bacteria treated with PR-26 were more susceptible to neutrophil phagocytosis (Fig. 8; PR-26 was different from control, $P < 0.05$). A 10 min. exposure of *S. choleraesuis* to 2 MICs of PR-26 significantly increased the capability of porcine neutrophils to phagocytose the bacteria. Treatment of *S. choleraesuis* with PR-39 at 8 MICs did not increase the phagocytic index of porcine neutrophils. Neutrophils phagocytosed both single and filamentous bacteria. These data show that, in addition to the direct antibacterial activity of PR-26, this antibacterial peptide predisposes enteric bacteria to elimination by phagocytic cells. This property suggests that PR-26 works synergistically with the host's immune system to limit enteric bacterial growth.

15 Influence of PR-39 on neutrophil chemotaxis. Phagocytic cells migrate from the blood to areas of inflammation in response to chemotactic agents. Fig. 9 shows that PR-14, PR-15, PR-16, PR-26, and PR-39 are chemotactic agents for neutrophils (PR-14, PR-15, PR-16, and PR-26 are used at 1 μM , and PR-39 was used at 0.05 μM ; the chemoattractant C5a, a positive control, was used at $1 \times 10^{-8}\text{M}$; starred entries are different from the control, $P < 0.05$). Fig. 10 shows a dose response of PR-39 for neutrophil chemotaxis. The ability of PR-39 to function as a chemotactic agent increases the probability that sufficient phagocytic cells are present at an inflammatory site to limit an infection.

25 Influence of PR-26 and PR-39 on intestinal epithelial cells. Fig. 11 shows the cytotoxic activity of PR-26 and PR-39 on rat small-intestine epithelial cells (IEC-6). PR-26 was not cytotoxic to IEC-6 cells even at concentrations (20 μM) much greater than the MIC for this peptide. However, IEC-6 cells were sensitive to PR-39 as cytotoxicity occurred at 0.5 μM , which is lower than the MIC for this peptide. These data show that PR-26 does not damage cells of the small intestine and should, therefore, be a safe oral antibiotic.

30 Effect of PR-26 on survival of mice challenged with *S. typhimurium*. In the first experiment, mice that received PR-26 were resistant to *Salmonella* (0 and 1 death for 250 and 100 $\mu\text{g}/\text{mouse}$, respectively), but the control group had a 27% mortality rate (Fig. 12; both PR-26 treatments were significantly different from controls, $P < 0.05$). Based on these data, PR-26 was effective in increasing the survival rates in this *Salmonella*-resistant strain of mice. There

35

-15-

was no evidence of toxicity caused by PR-26. In the second experiment, both 50 and 100 µg PR-26 were sufficient to increase survival to more than 50% at 14 days post-infection (Fig. 13; both PR-26 treatments were different from the PBS control at days 6 through 14, $P < 0.05$, but not different from each other. In the third experiment, administration of PR-26 was most effective when given at time 0; however, the 24-hour post-infection treatment slightly improved the survival compared to the PBS treatment by day 14 (Fig. 14; treatments at 0, 24, and 72 hrs. were different from PBS control at day 15, $P < 0.05$ for treatments at 0 and 24 hrs., $P < 0.10$ for treatment at 72 hrs.). The delivery of PR-26 at 48 hrs. post-infection was not different than the PBS control treatment. The rapidity of uptake of *S. typhimurium* and delivery to the liver and spleen during an i.p. infection may be the cause of the reduced effectiveness of PR-26 by 48 hrs. post-infection.

Regulation of neutrophil O_2^- production by PR-26 and PR-39.

(i) O_2^- production assays. When PR-39 was incubated with neutrophils for at least 45 min. before stimulation with phorbol myristate acetate (PMA), a significant reduction in O_2^- generation was observed (Fig. 15; neutrophils (1×10^6) were incubated without (Control) or with PR-39 (5 µM) for the indicated times before stimulation with PMA; values are means \pm SEM, $n=2$; starred entries are different from control, $P < 0.05$). This effect was even more apparent with longer pre-incubation periods, approaching as much as 70% oxidase inhibition. PR-26 also significantly reduced O_2^- generation (Fig. 16; neutrophils were incubated with PR-39 or derivatives for 150 min. before stimulation with PMA; results are reported as % relative to control (no peptides); however, the statistical analysis was based on nmoles of O_2^- produced; values are means, $n=3$; starred entries are different from control, $P < 0.05$; this experiment was conducted twice with similar results). However, the shorter peptides, PR-14, PR-15, PR-16, PR-19, and PR-23, did not significantly reduce O_2^- generation by intact neutrophils. PR-39 (20 µM) added to neutrophils at the same time as PMA activities did not affect generation of O_2^- , while longer incubation periods with this peptide did not affect neutrophil viability, as judged by trypan blue dye exclusion.

To investigate the inhibition of neutrophil O_2^- generation by PR-39 and its fragments in more detail, a cell-free assay system of O_2^- generation was used. PR-26 and PR-39 were very potent inhibitors of O_2^- generation in the

-16-

cell-free assay; concentrations that inhibited 50% (IC_{50}) of O_2^- generation were approximately 1 and 2 μM , respectively (Fig. 17; data are the average of duplicate reactions and are representative of 3 to 4 independent experiments; control (100%) activities ranged from 0.9 to 2.4 nmol O_2^- /min/ 5×10^5 cell equivalents of membrane). At concentrations greater than 5 μM , both proline-rich peptides completely inhibited the generation of O_2^- . PR-15 and PR-16 did not affect O_2^- generation; however, PR-14 did reduce O_2^- generation at concentrations greater than 25 μM .

PR-19 and PR-23 had some oxidase inhibitory activity; IC_{50} 's of PR-19 and PR-23 were approximately 5 μM and 25 μM , respectively (Fig. 17). These findings suggest that both peptides contain the oxidase inhibitory domain, although the first three arginine residues of PR-26 contribute greatly to oxidase inhibitory activity.

(ii) Binding assays. The findings that PR-26 and PR-39 inhibited both cell-free and whole cell O_2^- production by neutrophils and that inhibition required preincubation of cells with peptides for at least 45 min. prior to PMA stimulation suggested that these peptides act through some intracellular target, such as the NADPH oxidase components themselves. To determine if PR-39 bound to specific neutrophil cytosol components, human and porcine cytosolic proteins were separated by SDS-PAGE, transferred to a PVDF membrane, and probed with biotinylated PR-39. Using this ligand-blot binding assay, it was found that PR-39 bound to a 47 kDa protein in both human and porcine cytosol preparations (Fig. 18; cytosolic protein (50 μg) from human and porcine polymorphonuclear (hPMN and pPMN) leukocytes was subjected to SDS-PAGE, transferred to a PVDF membrane, and then probed with biotinylated PR-39; molecular masses of standards in kDa are shown on the left of the blots).

Competitive binding assays were conducted to determine if PR-39 bound to recombinant $p47^{phox}$. These solution binding experiments with pure recombinant $p47^{phox}$ and PR-39 showed that increasing concentrations of nonbiotinylated PR-39 inhibited specific binding of biotinylated PR-39 to recombinant $p47^{phox}$ (Fig. 19; competitive binding analysis was conducted by incubating various concentrations of PR-39 with recombinant $p47^{phox}$ and then with biotinylated PR-39; after dialysis, solutions were dot-blotted onto PVDF membranes and probed for bound biotinylated PR-39). Binding was inhibited

-17-

90% at equimolar concentrations of labeled and unlabeled peptide (0.06 μ M) and was completely blocked by a tenfold excess of unlabeled peptide. This finding shows that PR-39 binds to p47^{phox}, that binding is specific, and implies that PR-39 decreases O₂⁻ generation by interfering with this cytosolic component of the NADPH oxidase complex.

At least two proline-rich sequences, the COOH-terminal region (residues 358-371) of p47^{phox} and the cytoplasmic region (residues 149-162) of p22^{phox}, are thought to bind to SH3 domains of p67^{phox} and p47^{phox}, respectively, and are essential for the activation of NADPH oxidase in vivo (Leto et al., 1994; Sumimoto et al., 1994; Finan et al., 1994; McPhail, 1994; de Mendez et al., 1996). Data obtained by probing neutrophil cytosolic proteins with biotinylated PR-39 suggested that PR-39 did not bind to p67^{phox} (Fig. 18). Because PR-39 bound directly to p47^{phox}, it was reasoned that PR-39 could interfere with assembly of NADPH oxidase by blocking its interaction with p22^{phox}.

To examine this hypothesis, solution binding assays were conducted between GST-p47^{phox} SH3 domain fusion protein and a recombinant p22^{phox} cytoplasmic domain (residues 127-195) in the presence or absence of PR-39 or its derived fragments. PR-26 and PR-39 effectively blocked the interaction between GST-p47^{phox} and recombinant p22^{phox} at concentrations close to their IC₅₀'s observed in the cell-free oxidase assay (Fig. 20; PR-26 or PR-39 were mixed with recombinant p22^{phox} and incubated for 60 min. with GST-p47^{phox}-SH3 fusion proteins (0.5 μ g) bound to glutathione-Sepharose beads and analyzed as described; after SDS-PAGE, Sepharose-bound proteins were analyzed by immunoblotting with anti-p22^{phox} antibody; controls included blotting of recombinant p22^{phox} lysate (5 x 10³ cell equivalents) alone (p22) and bound complexes detected without competing PR-peptides (-); concentrations of competing peptides defined above each blot are indicated below the blots). Using this binding assay, analysis of the inhibition by PR-14, PR-15, PR-16, PR-19, and PR-23 revealed the importance of the COOH-terminal region of PR-26 in SH3 binding (Fig. 21; inhibition by PR-19 or PR-23, conducted as in Fig. 20) (Fig. 22; comparison of inhibitory activities of smaller peptides at 25 μ M). These data not only supported the hypothesis that PR-39 blocks the interaction between p47^{phox} and p22^{phox}, but also indicated that the main structural motif involved in the interaction between PR-39 and SH3 domains of p47^{phox} was in the central segment (PR-16) of PR-39. This 16-residue segment

-18-

contains structural elements compatible with consensus features of both classes of SH3 peptide ligands (Feng et al., 1994; Lim et al., 1994).

Two sequences within PR-16, RPPPFPP (Sequence ID No. 8) and PPRLPPRI (Sequence ID No. 9), conform to consensus sequences for either Class I (+) and Class II (-) binding orientations, $X_1pX_2PpX_3P$ and $X_3PpX_2PpX_1$, respectively (upper-case denotes critical contact residues, P denotes proline, and X_1 or X_3 favor arginine residues). Since PR-23 inhibits better than PR-19, a Class II binding orientation appears likely. However, because PR-19 had greater inhibitory activity in the cell-free assay but was less potent in blocking the interaction of $p47^{phox}$ SH3 domains with $p22^{phox}$ when compared to PR-23, the polybasic motif of the amino-terminus of PR-26 also represents a separate important component of oxidase inhibition. This observation is supported by studies showing that several other cationic peptides effectively inhibit NADPH oxidase, including peptides derived from $p47^{phox}$ (Joseph et al., 1994). Thus, critical regions at both ends of PR-26 have been defined based on dramatic losses of function seen with the deleted forms, PR-19 and PR-23.

The polybasic motif at the amino-terminus of PR-26 and PR-39, in addition to their overall amphipathic character, may have a role in promoting internalization of these inhibitory peptides, since similar structural properties are thought to enable internalization of various synthetic peptides designed for intracellular targets (Derossi et al., 1994; Mann et al., 1994; Fawell et al., 1994; Vellette et al., 1994; Taffs et al., 1992). PR-39 uptake may involve an active endocytic process or direct membrane lipid interactions, since this peptide binds and induces conductance changes in pure lipid bilayers (Cabiaux et al., 1992).

PR-39 is the first naturally occurring down regulator of phagocyte NADPH oxidase identified that interferes with assembly of this enzyme by binding to $p47^{phox}$. Little is known about the mechanisms governing respiratory burst kinetics, although models based on a continuous cycling of cytosolic components are consistent with the direct inhibitory effects of PR-39 on oxidase component interactions. The paradoxical finding of a neutrophil peptide possessing both antibacterial and oxidase inhibitory activities is intriguing, since a switch from oxygen-dependent to oxygen-independent

-19-

bactericidal mechanisms by an accumulation of this peptide within inflammatory sites could serve several important functions.

PR-39, which has a well-documented role as an antibacterial peptide, might have several roles in tissue repair by directly inhibiting NADPH oxidase activity and limiting related proinflammatory responses, while also affecting gene expression patterns that promote wound healing (Gallo et al., 1994). Furthermore, since reactive oxygen intermediates have also been shown to function as second messengers that can regulate gene expression (Schreck et al., 1991), PR-39 may indirectly influence gene expression patterns related to oxidative stress. These opposing activities of PR-39 illustrate a fine balance required in host defense mechanisms: antibacterial activity is necessary to control microbial pathogens and oxidase inhibitory activity is important for restricting tissue damage caused by excessive oxygen radicals generated by NADPH oxidase. In the case of PR-39, one peptide possesses both activities. These findings suggest a mechanism for interaction between oxidative and nonoxidative antimicrobial systems of neutrophils and may serve as a basis for design of drugs effective against production of oxidants in chronic or acute inflammatory disease states.

-20-

REFERENCES

The following references are incorporated by reference herein.

- 5 Abo, A., Boyhan, A., West, I., Thrasher, A.J. & Segal, A.W. (1992) *J. Biol. Chem.* **267**, 16767-16770.
- Agerberth, B., Lee, J.Y., Bergman, T., Boman, H.G., Mutt, V. & Tornvall, H. (1991) *Eur. J. Biochem.* **202**, 849-854.
- 10 Boman, H.G. (1991) *Cell* **65**, 205-207.
- Boman, H.G. (1995) *Annu. Rev. Immunol.* **13**, 61-92.
- 15 Boman, H.G., Agerberth, B. & Boman, A. (1993) *Infect. Immun.* **61**, 2978-2984.
- Cabiaux, V., Agerberth, B., Johansson, J., Homblé, F., Goormaghtigh, E. & Ruyschaert, J.-M. (1994) *Eur. J. Biochem.*, **224**, 1019-1027.
- 20 DeLeo, F.R., Nauseef, W.M., Jesaitis, A.J., Burritt, J.B., Clark, R.A. & Quinn, M.T. (1995) *J. Biol. Chem.* **270**, 26246-26251.
- de Mendez, I., Adams, A.G., Sokolic, R.A., Malech, H.L. & Leto, T.L. (1996) *EMBO J.*, in press.
- 25 Demling, R.H. (1990) *Circ. Shock* **30**, 297-309.
- Derossi, D., Joliot, A.H., Chassaing, G. & Prochiantz, A. (1994) *J. Biol. Chem.* **269**, 10444-10450.
- 30 Fawell, S., Seery, J., Daikh, Y., Moore, C., Chen, L.L., Pepinsky, B. & Barsoum, J. (1994) *Proc. Natl. Acad. Sci. USA* **91**, 664-668.
- Feng, S., Chen, J.K., Yu, H., Simon, J.A. & Schreiber, S.L. (1994) *Science* **266**, 1241-1247.
- 35

-21-

Finan, P., Shimizu, Y., Gout, I., Hsuan, J., Truong, O., Butcher, C., Bennett, P.,
Waterfield, M.D. & Kellie, S. (1994) *J. Biol. Chem.* **269**, 13752-13755.

Gabay, J.E. (1994) *Science* **264**, 373-374.

5

Gallo, R.L., Ono, M., Povsic, T., Page, C., Eriksson, E., Klagsbrun, M. &
Bernfield, M. (1994) *Proc. Natl. Acad. Sci. USA* **91**, 11035-11039.

Gennaro, R., Skerlavaj, B. & Romeo, D. (1989) *Infect. Immun.* **57**, 3142-3146.

10

Granger, D.N. & Korthuis, R.J. (1995) *Annu. Rev. Physiol.* **57**, 311-332.

Hopp, T.P. (1985) in *Synthetic Peptides in Biology and Medicine*, Alitalo, K.,
Partanen, P. & Vaheri, A. (Elsevier Science Publishers), pp. 3-12.

15

Joseph, G., Gorzalczany, Y., Koshikin, V. & Pick, E. (1994) *J. Biol. Chem.* **269**,
29024-29031.

Kilpatrick, L.E., Jakobovics, E., McCawley, L.J., Kane, L.H. & Korchak, H.M.
(1995) *J. Immunol.* **154**, 3429-3436.

20

Klebanoff, S.J. (1992) in *Inflammation: Basic Principles and Clinical
Correlates*, eds. Gallin, J.I., Goldstein, I.M. & Snyderman, R. (Raven, New
York), pp. 541-588.

25

Lee, J.-Y., Boman, A., Chuanxin, S., Andersson, M., Jörnvall, H., Mutt, V. &
Boman, H.G. (1989) *Proc. Natl. Acad. Sci. USA* **86**, 9159-9162.

Lehrer, R.I., Lichtenstein, A.K. & Ganz, T. (1993) *Annu. Rev. Immunol.* **11**,
105-128.

30

Leto, T.L., Garrett, M.C., Fujii, H. & Nuno, H. (1991) *J. Biol. Chem.* **266**, 19812-
19818.

-22-

- Leto, T.L., Adams, A.G. & de Mendez, I. (1994) *Proc. Natl. Acad. Sci. USA* **91**, 10650-10654.
- 5 Lim, W.A., Richards, F.M. & Fox, R.O. (1994) *Nature*, **372**, 375-379.
- Litteri, L. & Romeo, D. (1993) *Infect. Immun.* **61**, 966-969.
- MacKenzie, F.M. & Gould, I.M. (1993) *J. Antimicrobial Chemotherapy* **32**, 519-537.
- 10 Malech, H.L. & Gallin, J.I. (1987) *New England J. Med.* **317**, 687-694.
- Maloy, W.L. & Prasad Kari, U. (1995) *Biopolymers (Peptide Science)* **37**, 105-122.
- 15 Mann, D.A. & Frankel, A.D. (1991) *EMBO J.* **10**, 1733-1739.
- Martin, E., Ganz, T. & Lehrer, R.I. (1995) *J. Leukoc. Biol.* **58**, 128-136.
- 20 Martinez-Cayuela, M. (1995) *Biochimie* **77**, 147-161.
- McPhail, L.C. (1994) *J. Exp. Med.* **180**, 2011-2015.
- 25 McPhail, L.C., Qualliotine-Mann, D., Agwu, D.E. & McCall, C.E. (1993) *Eur. J. Haematol.* **51**, 294-300.
- Moore, K.S., Bevins, C.L., Tomassini, N., Huttner, K.M., Sadler, K., Moreira, J.E., Reynolds, J. & Zasloff, M. (1992) *J. Histochem. Cytochem.* **40**, 367-378.
- 30 National Committee for Clinical Laboratory Standards. Methods for Dilution Antimicrobial Susceptibility Tests for Bacteria that Grow Aerobically, 2nd Ed.; Approved Standard NCCLS Document M7-A2, Vol 10, No. 8.

-23-

Rotrosen, D. (1992) in *Inflammation: Basic Principles and Clinical Correlates*, eds. Gallin, J.I., Goldstein, I.M. & Snyderman, R. (Raven, New York), pp. 589-601.

5 Rotrosen, D., Yeung, C.L., Leto, T.L., Malech, H.L. & Kwong, C.H. (1992) *Science* **256**, 1459-1462.

Salak, J.L., McGlone, J.J. & Lyte, M. (1993) *Vet. Immunol. Immunopathol.* **39**, 327-337.

10 Schreck, R., Rieber, P. & Baeuerle, P.A. (1991) *EMBO J.*, **10**, 2247-2258.

Selsted, M.E. & Ouellette, A.J. (1995) *Trends Cell Biol.* **5**, 114-119.

15 Shasby, D.M., Vanbenthuyssen, K.M., Tate, R.M., Shasby, S.S., McMurtry, I. & Repine, J.E. (1982) *Am. Rev. Resp. Dis.* **125**, 443-447.

Shi, J., Goodband, R.D., Chengappa, M.M., Nelssen, J.L., Tokach, M.D., McVey, D.S. & Blecha, F. (1994a) *J. Leukoc. Biol.* **56**, 88-94.

20 Shi, J., Ross, C.R., Chengappa, M.M. & Blecha, F. (1994b) *J. Leukoc. Biol.* **56**, 807-811.

Shi, J. & Blecha, F. (1995) *Conf. Res. Workers Anim. Dis.*, p. 48.

25 Shi, J., Ross, C.R., Chengappa, M.M., Sylte, M.J., McVey, D.S. & Blecha, F. (1996) *Antimicrob. Agents Chemother.*, **40**, 115-121.

Steiner, H., Hultmark, D., Engström, A. Bennich, H. & Boman, H.G. (1981) *Nature* **292**, 246-248.

30 Sumimoto, H., Kage, Y., Nunoi, H., Sasaki, H., Nose, T., Fukumaki, Y., Ohno, M., Minakami, S. & Takeshige, K. (1994) *Proc. Natl. Acad. Sci. USA* **91**, 5345-5349

35 Taffs, R.E. & Sitkovsky, M.V. (1992) *J. Pharmaceutical Sci.* **81**, 37-44.

-24-

Vallette, F.M., Juin, P., Pelleschi, M. & Henry, J.-P. (1994) *J. Biol. Chem.* **269**, 13367-13374.

Zasloff, M. (1987) *Proc. Natl. Acad. Sci. USA* **84**, 5449-5543.

5

Zasloff, M. (1992) *Current Opin. Immunol.* **4**, 3-7.

-25-

SEQUENCE LISTING

(1) GENERAL INFORMATION:

5

(i) APPLICANT: Blecha, Frank
Shi, Jishu

(ii) TITLE OF INVENTION: Synthetic Antimicrobial Peptide

10

(iii) NUMBER OF SEQUENCES: 7

(iv) CORRESPONDENCE ADDRESS:

(A) ADDRESSEE: Hovey, Williams, Timmons & Collins

15

(B) STREET: 2405 Grand Blvd., Ste. 400

(C) CITY: Kansas City

(D) STATE: Missouri

(E) COUNTRY: U.S.A.

(F) ZIP: 64108

20

(v) COMPUTER READABLE FORM:

(A) MEDIUM TYPE: Floppy disk

(B) COMPUTER: IBM PC compatible

(C) OPERATING SYSTEM: PC-DOS/MS-DOS

25

(D) SOFTWARE: PatentIn Release #1.0, Version #1.25

(vi) CURRENT APPLICATION DATA:

(A) APPLICATION NUMBER: 08/419,066

(B) FILING DATE: April 10, 1995

30

(C) CLASSIFICATION:

(viii) ATTORNEY/AGENT INFORMATION:

(A) NAME: Collins, John M.

(B) REGISTRATION NUMBER: 26,262

35

(C) REFERENCE/DOCKET NUMBER: 23625

(ix) TELECOMMUNICATION INFORMATION:

(A) TELEPHONE: (816) 474-9050

(B) TELEFAX: (816) 474-9057

40

-26-

(2) INFORMATION FOR SEQ ID NO:1:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 39 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

Arg Arg Arg Pro Arg Pro Pro Tyr Leu Pro Arg Pro Arg Pro Pro Pro
1 5 10 15

Phe Phe Pro Pro Arg Leu Pro Pro Arg Ile Pro Pro Gly Phe Pro Pro
20 25 30

Arg Phe Pro Pro Arg Phe Pro
35

(3) INFORMATION FOR SEQ ID NO:2:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 26 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

Arg Arg Arg Pro Arg Pro Pro Tyr Leu Pro Arg Pro Arg Pro Pro Pro
1 5 10 15

Phe Phe Pro Pro Arg Leu Pro Pro Arg Ile
20 25

-27-

(4) INFORMATION FOR SEQ ID NO:3:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 23 amino acids

5 (B) TYPE: amino acid

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

10 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

Pro Arg Pro Pro Tyr Leu Pro Arg Pro Arg Pro Pro Pro Phe Phe Pro
1 5 10 15

15 Pro Arg Leu Pro Pro Arg Ile
20

(5) INFORMATION FOR SEQ ID NO:4:

20

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 19 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

25

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

30 Arg Arg Arg Pro Arg Pro Pro Tyr Leu Pro Arg Pro Arg Pro Pro Pro
1 5 10 15

Phe Phe Pro

35

-28-

(6) INFORMATION FOR SEQ ID NO:5:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 16 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

Arg Pro Arg Pro Pro Pro Phe Phe Pro Pro Arg Leu Pro Pro Arg Ile
1 5 10 15

(7) INFORMATION FOR SEQ ID NO:6:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 15 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

Arg Ile Pro Pro Gly Phe Pro Pro Arg Phe Pro Pro Arg Phe Pro
1 5 10 15

(8) INFORMATION FOR SEQ ID NO:7:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 14 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

Arg Arg Arg Pro Arg Pro Pro Tyr Leu Pro Arg Pro Arg Pro
1 5 10

-29-

(9) INFORMATION FOR SEQ ID NO:8:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 7 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

Arg Pro Pro Pro Phe Phe Pro
1 5

(10) INFORMATION FOR SEQ ID NO:9:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 8 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:

Pro Pro Arg Leu Pro Pro Arg Ile
1 5

-30-

Claims:

1. A method of inhibiting leukocyte O_2^- production comprising the step of administering to a leukocyte a peptide capable of inhibiting leukocyte O_2^- production.
2. The method of claim 1, wherein the peptide has a sequence included in Sequence ID No. 1.
3. The method of claim 2, wherein the peptide has the sequence selected from the group consisting of Sequence ID Nos. 1 and 2.
4. The method of claim 1, wherein the peptide has a length of less than 60 amino acid residues.
5. The method of claim 1, wherein the peptide is synthesized.
6. The method of claim 1, wherein the peptide inhibits the activity of NADPH oxidase.
7. The method of claim 1, wherein the leukocyte is a mammalian leukocyte.
8. The method of claim 7, wherein the leukocyte is a porcine leukocyte.
9. The method of claim 1, wherein the leukocyte is a neutrophil.
10. A method of attracting a leukocyte to a location comprising the step of administering to the location a peptide capable of attracting a leukocyte.
11. The method of claim 10, wherein the peptide has a sequence included in Sequence ID No. 1.

-31-

12. The method of claim 11, wherein the peptide has the sequence selected from the group consisting of Sequence ID Nos. 1, 2, 5, 6, and 7.

5 13. The method of claim 10, wherein the peptide has a length of less than 60 amino acid residues.

10 14. The method of claim 10, wherein the peptide is synthesized.

15 15. The method of claim 10, wherein the leukocyte is a mammalian leukocyte.

15 16. The method of claim 15, wherein the leukocyte is a porcine leukocyte.

20 17. The method of claim 10, wherein the leukocyte is a neutrophil.

1/12







Fig. 1.

Fig. 2.




2/12

MW _{ca}	1813	1777	1939	3230	4720
MW _{sp}	1812	1777	1941	3230	4720

A*Fig. 3.***B***Fig. 4.*



3/12

	1	
2	3	4
5	6	7
8	9	10
11	12	13

*Fig. 5.**Fig. 6.*

4/12

Postantibiotic Effect of PR-39 and PR-26

*Fig. 7.*Phagocytic Susceptibility of Salmonella
after Exposure to PR-26 & PR-39*Fig. 8.*

5/12

Chemotaxis of Porcine Neutrophils by
PR-39 and It's Analogs




Fig. 9.

Cytotoxicity of PR-26 & PR-39
in IEC-6 Cells






Fig. 11.

6/12

*Fig. 10.*

Effect of PR-26 on an Oral Challenge
with *Salmonella typhimurium*

*Fig. 12.*

7/12

Survival of Mice Challenged i.p. with *Salmonella typhimurium*




Fig. 13.

Effect of PR-26 on an Oral Challenge with *Salmonella typhimurium*








Fig. 14.


8/12

*Fig. 15.**Fig. 16.*



9/12

*Fig. 17.**Fig. 18.*

10/12

*Fig. 19.*

11/12

*Fig. 20.**Fig. 21.*

12/12

(-) PR-14
PR-16
PR-15
PR-26



(-25 μ M-)

Fig. 22.

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US96/04674

A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) : A 61K 38/16

US CL : 514/2

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 514/2, 524/12, 514/13, 514/14

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

APS, CAS Online

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	US 5,118,665 A (L. R. PICKART) 02 JUNE, 1992 see entire document.	1-17

☐ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

* Special categories of cited documents:	* "I" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
* "A" document defining the general state of the art which is not considered to be of particular relevance	* "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
* "E" earlier document published on or after the international filing date	* "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
* "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	* "Z" document member of the same patent family
* "O" document referring to an oral disclosure, use, exhibition or other means	
* "P" document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search 09 JULY 1996	Date of mailing of the international search report 26 AUG 1996
Name and mailing address of the ISA/US Commissioner of Patents and Trademarks Box PCT Washington, D.C. 20231 Facsimile No. (703) 305-3230	Authorized officer DAVID LUKTON Telephone No. (703) 308-0196